

Modulator simulations with open boundary and focusing field

Jun Ma, Roman Samulyak, Kwangmin Yu

Department of Applied Mathematics and Statistics
Stony Brook University

2016.3.3

Numerical setting

- Domain
 - Transversal : $4\text{e-}3\text{m}$ ($14 \times \text{Debyelength}$) open boundary
 - Longitudinal : $2\text{e-}5\text{m}$ ($20 \times \text{Debyelength}$) periodic boundary
- Grid number : 20 per Debye length
- Electron
 - Gaussian in x, y
 - Uniform in z
- Ion
 - At center
 - At one σ

Linear focusing field


$$\vec{E}_0(x) = \frac{m_e}{e} \frac{\sigma^2}{r_0^2} (x - x_0)$$

G. Bell et al., VLASOV AND PIC SIMULATIONS OF A MODULATOR SECTION FOR COHERENT ELECTRON COOLING, Proceedings of 2011 Particle Accelerator Conference, New York, NY, USA, MOP067.


Implemented as

$$\vec{E} = k\vec{x}$$


Transversal number distribution, initial




Transversal number distribution, without focusing field




Transversal number distribution, with focusing field




Transversal velocity distribution, initial




Transversal velocity distribution, without focusing field




Transversal velocity distribution, with focusing field




Transversal velocity distribution, without focusing field




Longitudinal number distribution




Longitudinal velocity distribution




Transversal number distribution




Transversal velocity distribution




Longitudinal number distribution



Longitudinal velocity distribution



Transversal number distribution



Transversal velocity distribution

