

PHY 564

Advanced Accelerator Physics

Lecture 7

Vladimir N. Litvinenko
Yichao Jing
Gang Wang

Department of Physics & Astronomy, Stony Brook University
Collider-Accelerator Department, Brookhaven National Laboratory

Review of Linear Algebra

Gang Wang

Matrix: definition and properties

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22} & \dots & a_{2m} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nm} \end{pmatrix}$$

Addition: $A + B = C \Leftrightarrow a_{ij} + b_{ij} = c_{ij}$

Multiplied by a constant: $kA = B \Leftrightarrow ka_{ij} = b_{ij}$

Equality: $A = B \Leftrightarrow a_{ij} = b_{ij}$

Multiplication (inner product): $AB = C \Leftrightarrow \sum_k a_{ik} b_{kj} = c_{ij}$

$$(AB)C = A(BC), \quad A(B + C) = AB + AC$$

In general $AB \neq BA$

Multiplication demands that A has the same number of columns as B has rows.

Matrix: special cases I

- Diagonal matrix: $a_{ij} = 0$ for $i \neq j$

$$A = \begin{pmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & 0 \\ \dots & 0 & \dots & \dots \\ 0 & 0 & \dots & a_{nn} \end{pmatrix}$$

If A and B are both diagonal matrix, they are commutative:

$$AB = BA$$

- Identity matrix: $AI = IA = A$ for $\forall A$

$$I = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & 0 & \dots & \dots \\ 0 & 0 & \dots & 1 \end{pmatrix}$$

$$I_{ij} = \delta_{ij} = \begin{cases} 1 & \text{for } i = j \\ 0 & \text{for } i \neq j \end{cases}$$

Matrix: special cases II

- Block diagonal matrix: A and A_i are square matrix.

$$A = \begin{bmatrix} A_1 & O & \dots & O \\ O & A_2 & \dots & O \\ \vdots & \vdots & \ddots & \vdots \\ O & O & \dots & A_k \end{bmatrix}, \quad A = \left[\begin{array}{ccc|c|cc} 1 & 3 & 2 & 0 & 0 & 0 \\ 7 & 0 & 2 & 0 & 0 & 0 \\ 1 & 1 & 2 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 6 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 3 & 3 \end{array} \right]$$

- Triangular matrix:

Upper diagonal matrix: elements below diagonal are all zero

$$U = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ 0 & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_{nn} \end{pmatrix}$$

Lower diagonal matrix: elements above diagonal are all zero

$$L = \begin{pmatrix} a_{11} & 0 & \dots & 0 \\ a_{21} & a_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix}$$

Matrix V: transpose matrix

- A matrix, B, is called the **transpose matrix** of a matrix A if

$$A_{ij} = B_{ji}$$

The transpose matrix is often denoted as A^T ,
i.e. $A_{ij} = (A^T)_{ji}$

- A square matrix A is called an **orthogonal matrix** if

$$A^T = A^{-1}$$

- A square matrix A is called **symmetric matrix** if

$$A^T = A \quad \text{and} \quad \text{anti-symmetric if} \quad A^T = -A$$

Matrix: trace

- In any square matrix, the sum of the diagonal elements is called the trace.

$$Tr(A) = \sum_i a_{ii}$$

- A useful property: $Tr(AB) = Tr(BA)$
- In general, $Tr(ABC) = Tr(BCA) \neq Tr(BAC)$
- Trace is a linear operator:

$$Tr(A + kB) = Tr(A) + k \cdot Tr(B)$$

Matrix: determinant of a matrix

- For a square matrix,

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix}$$

- The determinant

$$D = \begin{vmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{vmatrix} = \det(A)$$

is called the determinant of matrix A and is denoted by $\det(A)$.

Determinant I

$$D = \begin{vmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{vmatrix} \begin{matrix} \text{n columns} \\ \text{n rows} \end{matrix} = \sum_{i,j,k} \epsilon_{ijk\dots} a_{1i} a_{2j} a_{3k} \dots$$

$\epsilon_{ijk\dots}$ is Levi-Civita symbol

$$\epsilon_{ijk\dots} = \begin{cases} 1 & \text{if } (i,j,k\dots) \text{ is even permutation of } (1, 2, 3\dots) \\ -1 & \text{if } (i,j,k\dots) \text{ is odd permutation of } (1, 2, 3\dots) \\ 0 & \text{if any of the two indices is repeated} \end{cases}$$

$$\epsilon_{ijk\dots l\dots m\dots} = -\epsilon_{ijk\dots m\dots l\dots}$$

Determinant II

A determinant of n dimension can be expanded over a column (or a row) into a sum of n determinants of $n-1$ dimension:

$$D = \begin{vmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & \dots & a_{2j} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ a_{n1} & \dots & a_{nj} & \dots & \dots \end{vmatrix} = \sum_i C_{ij} a_{ij}$$

$$M_{ij} = \begin{vmatrix} \overbrace{a_{11} \dots a_{1j} \dots a_{1n}}^{n-1 \text{ columns}} \\ \dots \\ \underbrace{a_{i1} \dots a_{ij} \dots a_{in}}_{n-1 \text{ rows}} \\ \dots \\ a_{n1} \dots a_{nj} \dots a_{nn} \end{vmatrix}$$

$C_{ij} = (-1)^{i+j} M_{ij}$ is called the ij^{th} cofactor of D .

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{21} \begin{vmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{vmatrix} + a_{31} \begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix}$$

Determinant III

- Multiplied by a constant

$$\begin{vmatrix} ka_{11} & a_{12} & a_{13} \\ ka_{21} & a_{22} & a_{23} \\ ka_{31} & a_{32} & a_{33} \end{vmatrix} = k \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

- The value of a determinant is unchanged if a multiple of one column (row) is added to another column (row)

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} a_{11} + ka_{12} & a_{12} & a_{13} \\ a_{21} + ka_{22} & a_{22} & a_{23} \\ a_{31} + ka_{32} & a_{32} & a_{33} \end{vmatrix}$$

- A determinant is equal to zero if any two columns (rows) are proportional

$$\begin{vmatrix} a_{12} & ka_{12} & a_{13} \\ a_{22} & ka_{22} & a_{23} \\ a_{32} & ka_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} 0 & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & a_{32} & a_{33} \end{vmatrix} = 0$$

Linear equation system

- Existence of non-trivial solution of homogeneous equations

$$\begin{cases} a_{11}x + a_{12}y + a_{13}z = 0 \\ a_{21}x + a_{22}y + a_{23}z = 0 \\ a_{31}x + a_{32}y + a_{33}z = 0 \end{cases}$$

$$x \cdot D \equiv x \cdot \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} a_{11}x + a_{12}y + a_{13}z & a_{12} & a_{13} \\ a_{21}x + a_{22}y + a_{23}z & a_{22} & a_{23} \\ a_{31}x + a_{32}y + a_{33}z & a_{32} & a_{33} \end{vmatrix} = 0$$

Similarly, $y \cdot D = 0$ and $z \cdot D = 0$

- Thus a set of homogenous linear equations have non-trivial solutions only if the determinant of the coefficients, D , vanishes.

Matrix: properties of the determinant of a matrix

- Some properties of the determinant of matrices

- $\det(A^T) = \det(A)$

- $\det(kA) = k^n \det(A)$

- $\det(AB) = \det(A)\det(B)$

Proof of $\det(AB) = \det(A)\det(B)$:

$$\begin{aligned}
 (1) \quad & \sum_{i,j,k \dots} \epsilon_{ijk \dots} a_{\beta i} a_{\alpha j} a_{\gamma k} \dots \\
 &= \sum_{j,i,k \dots} \epsilon_{jik \dots} a_{\beta j} a_{\alpha i} a_{\gamma k} \dots \\
 &= \sum_{i,j,k \dots} \epsilon_{jik \dots} a_{\alpha i} a_{\beta j} a_{\gamma k} \dots \\
 &= - \sum_{i,j,k \dots} \epsilon_{ijk \dots} a_{\alpha i} a_{\beta j} a_{\gamma k} \dots
 \end{aligned}$$

$$\begin{aligned}
 (3) \quad & |AB| = \sum_{i,j,k \dots} \epsilon_{ijk \dots} (AB)_{1i} (AB)_{2j} (AB)_{3k} \dots \\
 &= \sum_{i,j,k \dots} \sum_{\alpha, \beta, \gamma} \epsilon_{ijk \dots} A_{1\alpha} B_{\alpha i} A_{2\beta} B_{\beta j} A_{2\gamma} B_{\gamma j} \dots \\
 &= \sum_{\alpha, \beta, \gamma} A_{1\alpha} A_{2\beta} A_{2\gamma} \dots \left\{ \sum_{i,j,k \dots} \epsilon_{ijk \dots} B_{\alpha i} B_{\beta j} B_{\gamma j} \dots \right\} \\
 &= |B| \sum_{\alpha, \beta, \gamma} \epsilon_{\alpha\beta\gamma \dots} A_{1\alpha} A_{2\beta} A_{2\gamma} \dots \\
 &= |A| |B|
 \end{aligned}$$

$$\begin{aligned}
 (2) \quad & \sum_{i,j,k \dots} \epsilon_{ijk \dots} a_{\alpha i} a_{\beta j} a_{\gamma k} \dots \\
 &= \epsilon_{\alpha\beta\gamma \dots} \sum_{i,j,k \dots} \epsilon_{ijk \dots} a_{1i} a_{2j} a_{3k} \dots \\
 &= \epsilon_{\alpha\beta\gamma \dots} |A|
 \end{aligned}$$

Matrix IV: inversion

- Inversion of a square matrix A is to find a square matrix B such that

$$AB = BA = I$$

B is called the **inverse matrix** of A and often denoted by A^{-1} , i.e.

$$AA^{-1} = A^{-1}A = I$$

- One way to find the inverse matrix is by

$$(A^{-1})_{ij} = \frac{C_{ji}}{|A|}, \quad \text{where } C_{ji} \text{ is the } ji^{th} \text{ cofactor of } A$$

Matrix VI: similarity transformation and diagonalization

- Two matrix, A and B, are called similar if there exists a invertible matrix P such that

$$B = P^{-1}AP,$$

and the transformation from A to B is called **similarity transformation**.

- **Diagnolization** of a matrix, A, is to find a similarity transformation matrix, P, such that $P^{-1}AP$ is a diagonal matrix:

Matrix VII: diagonalization

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \lambda_n \end{pmatrix} \quad \text{or} \quad AP = P \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \lambda_n \end{pmatrix}$$

- If we look at the j^{th} column of the second equation, it follows

$$\sum_k a_{ik} p_{kj} = \sum_k p_{ik} \lambda_k \delta_{kj} = \lambda_j p_{ij} \quad (*)$$

Defining a $n \times 1$ matrix (i.e. a column vector) $|P^j\rangle$

such that $|P^j\rangle_i = p_{ij}$ (note: j is fixed)

Equation (*) becomes: $A|P^j\rangle = \lambda_j |P^j\rangle$

Matrix VIII: eigenvalue and eigenvector

- For a matrix A , a vector matrix X is called an **eigenvector** of A if

$$A \cdot X = \lambda X$$

where λ is called the **eigenvalue** associated with the eigenvector X .

- The eigenvalues are found by solving the following polynomial equation

$$(A - \lambda I) \cdot X = 0 \Rightarrow \det(A - \lambda I) = 0$$

Defective Matrix

- Not all square matrix can be diagonalized:

$$A = \begin{pmatrix} 2 & -3 \\ 3 & -4 \end{pmatrix}; \quad \det(A - \lambda I) = \begin{vmatrix} 2 - \lambda & -3 \\ 3 & -4 - \lambda \end{vmatrix} = 0 \Rightarrow (\lambda + 1)^2 = 0 \Rightarrow \lambda = -1$$

$$\begin{pmatrix} 2 & -3 \\ 3 & -4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = - \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \Rightarrow \begin{cases} 3x_1 - 3x_2 = 0 \\ 3x_1 - 3x_2 = 0 \end{cases} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix};$$

We end up with only one eigenvector.

- A square matrix that does not have a complete set of eigenvectors is not diagonalizable and is called a **defective matrix**.
- If a matrix, A , is defective (and hence is not similar to a diagonal matrix), then what is the simplest matrix that A is similar to?

Jordan form matrix

- Definition: a **Jordan block** with value λ is a square, upper triangular matrix whose entries are all λ on the diagonal, all 1 on the entries immediately above the diagonal, and zero elsewhere:

$$J(\lambda) = \begin{bmatrix} \lambda & 1 & 0 & \cdots & 0 & 0 \\ 0 & \lambda & 1 & \cdots & 0 & 0 \\ 0 & 0 & \lambda & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda & 1 \\ 0 & 0 & 0 & \cdots & 0 & \lambda \end{bmatrix}$$

1D: $[\lambda]$

2D: $\begin{bmatrix} \lambda & 1 \\ 0 & \lambda \end{bmatrix}$

3D: $\begin{bmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 1 \\ 0 & 0 & \lambda \end{bmatrix}$

- Definition: a Jordan form matrix is a block diagonal matrix whose blocks are all Jordan blocks

- Theorem: Let A be a $n \times n$ matrix. Then there is a Jordan form matrix that is similar to A .

$$\begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 3 & 1 & 0 & 0 \\ 0 & 0 & 0 & 3 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{bmatrix}$$

Symplectic Matrix

$$M^T S M = S \quad (**)$$

$$S = \begin{pmatrix} S_{1D} & 0 & \dots & 0 \\ 0 & S_{1D} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & S_{1D} \end{pmatrix}; \quad S_{1D} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

- A matrix satisfying condition (**) is called a **symplectic matrix**.
 - inverse: $M^T S M = S \Rightarrow S M^T S M = S^2 = -I \Rightarrow (-S M^T S) M = I \Rightarrow M^{-1} = -S M^T S$
 - if M and N are both symplectic, then their product, MN, is also symplectic $(MN)^T S (MN) = N^T M^T S M N = N^T S N = S$
 - if M is symplectic, M^T is also symplectic $(M^T S M)^{-1} = -S \Rightarrow M^{-1} S (M^T)^{-1} = S \Rightarrow S = M S M^T \Rightarrow (M^T)^T S M^T = S$

Symplectic Matrix II

- If λ is eigen value then $1/\lambda$ is also an eigenvalue and the multiplicity of λ and $1/\lambda$ is the same.
 - It implies that the eigenvalues are coming in pairs $\{\lambda, 1/\lambda\}$.
- As a consequence of above property, the determinant of a symplectic matrix is 1.