

**PHY 554. Homework 2.**

*Handed: August 30*


*Return by: September 6*

*Electronic copies accepted at [vladimir.litvinenko@stonybrook.edu](mailto:vladimir.litvinenko@stonybrook.edu)*

**HW 1 (5 point):** Let's first determine an effective focal length,  $F$ , of the of a paraxial (e.g. small angles!) focusing object (a black-box) as ratio between a parallel displacement of trajectory at its entrance to corresponding change of the angle at its exit (see figure below):

$$F = -\frac{x}{x'}; x' \equiv \frac{dx}{dz}$$

see figure below for



For completeness, the distance from the entrance to the object to the trajectory crossing the axis,  $l$ , in general is not equal to the focal length. In beam optics this is frequently, but not correctly, referred as astigmatism – in contrast, the astigmatism is defined as dependence of the focal strengths on the direction of propagation of the ray (particle).

Let consider a doublet of two thin lenses: a focusing ( $F$ ) and defocusing ( $D$ ) lenses with equal but opposite in sign focal length  $F$  with center separated by distance  $L$  as in Fig. 1.





Fig.1. Two combinations of a doublet:  $FD$  and  $DF$ .

1. (3 points) Show through a calculation of the ray trajectory that the focal lengths of  $FD$  and  $DF$  doublets are equal and given by following expression:

$$F_{eff} = \frac{F^2}{L}$$

2. (2 points) Determine location of the ray crossing the axis and find their difference between  $FD$  and  $DF$  doublets – this indeed would be an astigmatism of doublet built from two quadruples.

*P.S. Definition (picture) of thin lens:*



**HW 2 (2 points):** Spectral brightness (sometimes called brilliance) of a light source is defined as

$$B = \frac{dN_{ph}}{dt d\Omega dA (d\lambda / \lambda)} = \frac{dN_{ph}}{dt d\Omega dA (d\omega / \omega)};$$

where  $\frac{dN_{ph}}{dt}$  is the number of photons per second with the spectral bandwidth  $d\omega / \omega$  radiated from an area  $dA$  into the solid angle  $d\Omega$ . The units used for brightness are expressed in photons per second

$$[B] = \frac{\text{photons}}{\text{sec} \cdot \text{mm}^2 \cdot \text{mrad}^2 (10^{-3} d\lambda / \lambda)}$$

As an exercise, calculate spectral brightness of NdYAG laser with average power of 10 W, wavelength of  $\lambda = 1.064 \mu\text{m}$ , Bandwidth of  $\Delta\omega = 700 \text{ GHz}$  and with diffraction limited spot size and angular spread:

$$\Delta x \cdot \Delta\theta_x = \frac{\lambda}{4\pi}; \Delta y \cdot \Delta\theta_y = \frac{\lambda}{4\pi}.$$

**HW 3 (3 points):** In a fixed Cartesian coordinates for a trajectory with  $\frac{dz}{dt} \neq 0$  of a particle moving in magnetic field  $\vec{B} = \hat{x}B_x + \hat{y}B_y + \hat{z}B_z$  equation for its trajectory can be written in terms of  $z$  as independent variable:

$$\frac{d^2x}{dz^2} = \frac{e}{p} \sqrt{1+x'^2+y'^2} (y'B_z - (1+x'^2)B_y + x'y'B_x);$$

$$\frac{d^2y}{dz^2} = -\frac{e}{p} \sqrt{1+x'^2+y'^2} (x'B_z - (1+y'^2)B_x + x'y'B_y);$$

$$x' \equiv \frac{dx}{dz}; y' \equiv \frac{dy}{dz};$$

where  $e$  is the particle's charge and  $p = \gamma mv$  is its relativistic momentum.

*Hint: consider constants of motion in a magnetic field.*