

## Homework 6.

### Problem 1. 3 x 5 points. Function of a Jordan block

(a) Show that powers of  $m \times m$  Jordan block

$$\mathbf{G} = \begin{bmatrix} \lambda & 1 & 0 & 0 \\ 0 & \lambda & 1 & 0 \\ 0 & 0 & \dots & 1 \\ 0 & 0 & 0 & \lambda \end{bmatrix}$$

are

$$\mathbf{G}^n = \begin{bmatrix} \lambda^n & C_1^n \lambda^{n-1} & C_2^n \lambda^{n-2} & \dots & C_k^n \lambda^{n-k} & C_{k+1}^n \lambda^{n-k-1} & \dots \\ 0 & \lambda^n & C_1^n \lambda^{n-1} & \dots & C_{k-1}^n \lambda^{n+1-k} & C_k^n \lambda^{n-k} & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \lambda^n \end{bmatrix}; C_k^n = \frac{n!}{(n-k)!k!} \quad (1)$$

Suggestion: use  $\mathbf{G}^0 = \mathbf{I}; \mathbf{G}^1 = \mathbf{G}$  - as first step, they satisfy (1). Then use induction assuming that (1) is correct for  $n$  and show that  $\mathbf{G}^{n+1} = \mathbf{G} \cdot \mathbf{G}^n$  satisfy (1) for  $n+1$ . Use a well know ratio  $C_k^{n+1} = C_k^n + C_{k-1}^n$ .

(b) For a polynomial function  $f(x) = \sum_{n=0}^N f_n x^n$  show that

$$f(\mathbf{G}) = \sum_{n=0}^N f_n \mathbf{G}^n = \begin{bmatrix} \sum_{n=0}^{\infty} f_n \lambda^n & \dots & \sum_{n=0}^{\infty} f_n C_k^n \lambda^{n-k} & \dots & \dots \\ 0 & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \sum_{n=0}^{\infty} f_n \lambda^n & \dots \end{bmatrix}$$

and  $\sum_{n=0}^N f_n C_k^n \lambda^{n-k} = \frac{1}{k!} \frac{d^k f}{d\lambda^k} \#$

(c) Prove that for an arbitrary (well behaved function!)  $f(x) = \sum_{n=0}^{\infty} f_n x^n$

$$(d) f(\mathbf{G}) = \begin{bmatrix} f(\lambda) & f'(\lambda)/1! & \dots & f^{(k)}(\lambda)/k! & \dots & f^{(n-1)}(\lambda)/(n-1)! \\ 0 & f(\lambda) & \dots & \dots & \dots & f^{(n-2)}(\lambda)/(n-2)! \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \dots & \dots & f'(\lambda)/1! \\ 0 & 0 & \dots & \dots & \dots & f(\lambda) \end{bmatrix}$$